

together with

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

Please see Important Safety Information for HIBSAGO on pages 3 and 4. Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

together with HIBSAGO (bepirovirsen)

is support that will **stay by your side** throughout your treatment.

Once your doctor has prescribed HIBSAGO, we will help you explore cost-saving options and keep you informed at every step.

It's part of our promise to understand your needs and help you move forward with confidence.

You're not alone. Let's do this together.

Please see Important Safety Information for HIBSAGO on pages 3 and 4. Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

Important Safety Information

What is HIBSAGO?

HIBSAGO is a prescription medicine used with nucleos(t)ide analog reverse transcriptase inhibitors (NrtIs) to treat adults with chronic hepatitis B virus (HBV) infection who do not have cirrhosis (advanced liver disease), have taken an NrtI for at least 6 months, have undetectable HBV DNA, and have an HBV surface antigen (HBsAg) level of 3,000 IU/mL or less.

What is the most important information I should know about HIBSAGO?

HIBSAGO may cause serious side effects, including:

- **flare-ups in liver enzyme levels during treatment** measured using liver function blood tests. **Tell your healthcare provider right away if you get any of these symptoms:**
  - upper right stomach pain
  - dark or "tea-colored" urine
  - yellowing of your skin or the white part of your eyes
  - nausea or vomiting
  - loss of appetite

- **low platelet counts (thrombocytopenia), which** can happen at any time during treatment, suddenly and without warning, and can be life-threatening. **Stop using HIBSAGO and tell your healthcare provider right away if you get any of these symptoms:**
  - unusual bruising or a rash of tiny reddish-purple spots
  - bleeding from skin cuts that does not stop or oozes
  - bleeding from your gums or nose
  - blood in your urine or stools
  - abnormal or heavy periods (menstrual bleeding)
  - sudden severe headaches or neck stiffness
  - bleeding into the whites of your eyes

Your healthcare provider will check your liver and platelet count with blood tests before and during treatment, and for up to 24 weeks after you stop HIBSAGO if results are abnormal.

Who should not use HIBSAGO?

Do not use HIBSAGO if you have:

- a condition where your liver is currently not working as well as it should
- advanced liver disease (cirrhosis)
- low platelet counts.

Before using HIBSAGO, tell your healthcare provider about all of your medical conditions, including if you:

- have or had bleeding problems

Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

Important Safety Information (cont'd)

- have kidney problems
- have received an organ transplant
- have an immune system problem (autoimmune disease) or a history of blood vessel inflammation (vasculitis)
- have hepatitis D. HIBSAGO is not recommended if you have hepatitis D because it may increase the chance of liver enzyme flare-ups.
- are pregnant or plan to become pregnant. It is not known if HIBSAGO will harm your unborn baby.
- are breastfeeding or plan to breastfeed. HIBSAGO may pass into your breast milk.

Tell your healthcare provider about all the medicines you take, especially:

- medicines that prevent blood clotting or lower your platelet count.
- medicines that may lower your kidney function.
- peginterferon alfa-2a.

What are the possible side effects of HIBSAGO?

HIBSAGO may cause these additional serious side effects, including:

- **Worsening kidney function.** Your healthcare provider will check your kidneys with blood tests before and during treatment, and for up to 12 weeks after you stop HIBSAGO if results are abnormal.
- **Inflammatory or immune system problems.** Tell your healthcare provider if you get flu-like symptoms; purple rashes or spots on legs; scaly, raised, or red patches; swelling and itchiness; or joint pain that does not go away.
- **Serious side effects when HIBSAGO is not injected correctly.** Rotate injection sites each week, keep the two weekly injections at least 2 inches (5 cm) apart, and do not inject into skin that is tender, bruised, red, or hard.
- **The most common side effects of HIBSAGO include:** injection site reactions, fever, tiredness, and headache.

These are not all the possible side effects of HIBSAGO. Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Please see accompanying full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

Here's what to expect from Together with HIBSAGO (bepirovirsen)



Coverage support

We'll check your insurance coverage and help you understand your benefits.*



Cost support

We can help you determine your eligibility for additional programs that may help you pay for HIBSAGO.



Patient support

We offer 1-to-1 support, education, and resources throughout your treatment.

Patient support is available in multiple languages for non-English speakers.

*The information provided by Together with HIBSAGO is not a guarantee of coverage or payment.

Please see Important Safety Information for HIBSAGO on pages 3 and 4. Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

Say hello to support at every step



Starting a new treatment may feel overwhelming, but you don't have to navigate it alone.

Whenever you have a question or concern, Together with HIBSAGO is only a phone call or message away.

- Copay assistance program
- Financial assistance
- Coverage questions
- Ongoing support

Please see Important Safety Information for HIBSAGO on pages 3 and 4. Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

How to enroll

Together with HIBSAGO offers a range of assistance options at no cost to you, but first, you and your doctor need to complete enrollment.

- Scan the QR code on the back of this brochure or visit TogetherwithHIBSAGO.com to begin.
- Once you enroll, you'll receive a call to guide you through next steps.

Have a quick question about HIBSAGO?
We're here to help.
1-8-HIBSAGO-11 (1-844-272-4611)

Only interested in copay?

We understand that cost matters. That's why we offer a copay assistance program to help make your treatment more affordable.

For Commercially Insured Patients

You could get HIBSAGO for as little as

$0*

per dose

The HIBSAGO Copay Program helps eligible commercially insured patients with their eligible out-of-pocket costs for HIBSAGO.

*Program annual maximums apply.
Visit TogetherwithHIBSAGO.com/costsupport for information about eligibility and full program terms and conditions.

Please see Important Safety Information for HIBSAGO on pages 3 and 4. Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

HIBSAGO (bepirovirsen)
Subcutaneous Use 150 mg/mL

Let's do this together

1-8-HIBSAGO-11 (1-844-272-4611)
Monday–Friday: 8 AM–8 PM ET

TogetherwithHIBSAGO.com

together with | HIBSAGO (bepirovirsen)

Please see Important Safety Information for HIBSAGO on pages 3 and 4. Please see full Prescribing Information, including Boxed Warning and Medication Guide, for HIBSAGO.

Trademarks are owned by or licensed to the GSK group of companies.

©2026 GSK or licensor.
PMUS-BPRBROC260029 October 2026
Produced in USA. 0002-0050-58

GSK