



# support

made just for you

tōgether

with

  
Zejula  
niraparib  
tablets 100/200/300 mg

Please see Important Safety Information on pages 3 and 4 and full [Prescribing Information](#), including [Patient Information](#), for ZEJULA at [ZEJULA.com](#).

together

with

  
Zejula  
niraparib  
tablets 100/200/300 mg

It's more than a program—it's **our promise** to understand **your needs**, listen to what matters most to you, and create meaningful **connections** along the way.

Once your doctor has prescribed ZEJULA, we will help you to **understand** the medication coverage process, investigate **affordable options**, and **guide you** through your care.

## APPROVED USES

ZEJULA is a prescription medicine used to treat adults for:

- maintenance treatment of advanced ovarian cancer, fallopian tube cancer, or primary peritoneal cancer with a certain type of abnormal *BRCA* gene or a positive laboratory tumor test for genomic instability called homologous recombination deficiency (HRD). ZEJULA is used after the cancer has responded (complete or partial response) to treatment with first-line platinum-based chemotherapy.
- maintenance treatment of ovarian cancer, fallopian tube cancer, or primary peritoneal cancer with a certain type of inherited (germline) abnormal *BRCA* gene that comes back. ZEJULA is used after the cancer has responded (complete or partial response) to treatment with platinum-based chemotherapy. Your healthcare provider will perform a test to make sure that ZEJULA is right for you.

It is not known if ZEJULA is safe and effective in children.

## IMPORTANT SAFETY INFORMATION

**ZEJULA may cause serious side effects, including:**

**Bone marrow problems called Myelodysplastic Syndrome (MDS) or a type of blood cancer called Acute Myeloid Leukemia (AML).** Some people who have ovarian cancer and who have received previous treatment with chemotherapy or certain other medicines for their cancer have developed MDS or AML during treatment with ZEJULA. MDS or AML may lead to death.

**Symptoms of low blood cell counts (low red blood cells, low white blood cells, and low platelets)** are common during treatment with ZEJULA. They can be a sign of serious bone marrow problems, including MDS or AML. These symptoms may include the following:

- Weakness
- Feeling tired
- Weight loss
- Frequent infections
- Fever
- Shortness of breath
- Blood in urine or stool
- Bruising or bleeding more easily

Your doctor will do blood tests to check your blood cell counts before treatment with ZEJULA. You will be tested weekly for the first month of treatment with ZEJULA, monthly for the next 11 months of treatment, and as needed afterward.

**High blood pressure** is common during treatment with ZEJULA, and it can become serious. Your doctor will check your blood pressure and heart rate at least weekly for the first two months, then monthly for the first year, and as needed thereafter during your treatment with ZEJULA.

**Posterior reversible encephalopathy syndrome (PRES)** is a condition that affects the brain and may happen during treatment with ZEJULA. If you have headache, vision changes, confusion, or seizure, with or without high blood pressure, please contact your doctor.

**Please see additional Important Safety Information on page 4 and full [Prescribing Information](#), including [Patient Information](#), for ZEJULA at [ZEJULA.com](#).**

## IMPORTANT SAFETY INFORMATION (cont'd)

**Before starting to take ZEJULA, tell your doctor about all of your medical conditions, including if you:**

- Have heart problems
- Have liver problems
- Have high blood pressure
- Are pregnant or plan to become pregnant. ZEJULA can harm your unborn baby and may cause loss of pregnancy (miscarriage)
  - If you are able to become pregnant, your doctor should perform a pregnancy test before you start treatment with ZEJULA
  - If you are able to become pregnant, you should use effective birth control (contraception) during treatment with ZEJULA and for 6 months after taking the last dose of ZEJULA
  - You should tell your doctor right away if you become pregnant
- Are breastfeeding or plan to breastfeed
  - ZEJULA may harm your baby. You should not breastfeed your baby during treatment with ZEJULA and for 1 month after taking the last dose of ZEJULA

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**The most common side effects of ZEJULA include the following:**

- |                                 |  |
|---------------------------------|--|
| ◦ Nausea                        | ◦ Shortness of breath                          |
| ◦ Tiredness                     | ◦ Rash   |
| ◦ Constipation                  | ◦ Diarrhea                                     |
| ◦ Pain in your muscles and back | ◦ Cough  |
| ◦ Pain in the stomach area      | ◦ Dizziness                                    |
| ◦ Vomiting                      | ◦ Changes in the amount or color of your urine |
| ◦ Loss of appetite              | ◦ Urinary tract infection                      |
| ◦ Trouble sleeping              | ◦ Low levels of magnesium in the blood         |
| ◦ Headache                      |  |

If you have certain side effects, then your doctor may change your dose of ZEJULA, temporarily stop, or permanently stop treatment with ZEJULA.

These are not all the possible side effects of ZEJULA. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects.

**Please see the full Prescribing Information, including Patient Information for ZEJULA.**

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088. You may also report negative side effects to GSK at [gsk.public.reportum.com](http://gsk.public.reportum.com) or call 1-888-825-5249.**

# Here's what to expect from Together with ZEJULA (niraparib)



## Coverage support

We'll check your insurance coverage and help you understand your benefits.\*

\*The information provided by Together with ZEJULA is not a guarantee of coverage or payment.



## Cost support

Your dedicated Patient Navigator<sup>†</sup> can help you determine your eligibility for additional programs that may help you pay for ZEJULA.



## Patient support

Your Patient Navigator<sup>†</sup> is at the heart of our program, offering tailored support throughout your treatment journey.

<sup>†</sup>The Patient Navigator does not provide medical advice and is not a part of your doctor's care team. Please ask your doctor any questions that you might have about your disease and treatment with ZEJULA.

# Here's what to expect from our Patient Experience Center



<sup>†</sup>The information provided by Together with ZEPJULA is not a guarantee of coverage or payment.




<sup>§</sup>Independent foundations are not affiliated with GSK and are operated independently. Together with ZEPJULA cannot guarantee that you will be eligible for assistance.

Please see Important Safety Information on pages 3 and 4 and full Prescribing Information, including Patient Information, for ZEPJULA at [ZEPJULA.com](http://ZEPJULA.com).

together  
with

Zejula  
niraparib  
tablets 100/200/300 mg

Visit **TogetherwithZEJULA.com**  
or call **1-844-4GSK-ONC (844-447-5662)**  
and start taking advantage of the help  
our Patient Experience Center can provide:

-  **Answers to coverage and benefits questions**
-  **Options to help with medication costs for which you may be eligible**
-  **Ongoing support and education about ZEJULA**

### For commercially insured

The ZEJULA Copay Program helps eligible commercially insured patients with their out-of-pocket costs for ZEJULA.

\*Subject to program maximums.

See full Terms and Conditions at [TogetherwithZEJULA.com/costsupport](https://www.togetherwithzejula.com/costsupport).

You could get  
ZEJULA for as little as

**\$0\***

per dose

### No commercial insurance?

To see whether you may be eligible for other support options, call your Patient Navigator at **1-844-4GSK-ONC (844-447-5662)**.



## Connect with us.

**1-844-4GSK-ONC (844-447-5662)**  
Monday–Friday (8:00 AM – 8:00 PM ET)

Visit [TogetherwithZEJULA.com](https://TogetherwithZEJULA.com) for more details  
or scan the QR code below to get started.



together  
with

  
ZeJula  
niraparib  
tablets 100/200/300 mg

Please see Important Safety Information on pages 3 and 4 and full [Prescribing Information](#),  
including [Patient Information](#), for ZEJULA at [ZEJULA.com](https://ZEJULA.com).

Trademarks are owned by or licensed to the GSK group of companies.

©2026 GSK or licensor.

PMUS-NRPBROC250032 February 2026

Produced in USA. 0002-0045-13

File name: Together with ZEJULA Overview Brochure

**GSK**